

Anatomy of Flowering Plants

I. Select the correct answer from the following questions:

Question 1.

Intercalary meristem is derived from

- (a) Apical meristem
- (b) Protoderm
- (c) Calyptragen
- (d) Lateral meristem

▼ [Answer](#)

Answer: (a) Apical meristem.

Question 2.

Secondary meristem develops from

- (a) Apical meristem
- (b) Permanent tissue
- (c) Secondary tissue
- (d) Vascular combium

▼ [Answer](#)

Answer: (b) Permanent tissue

Question 3.

Cambium is considered to be a lateral meristem because it

- (a) Gives rise to lateral branches
- (b) Increase the girth of the plant
- (c) Increase both length and girth of plant
- (d) Increase the length of the plant.

▼ [Answer](#)

Answer: (b) Increase the girth of the plant

Question 4.

Quiscent centre is located in

- (a) Shoot apex
- (b) Root apex
- (c) Bud apex
- (d) Leaf apex

▼ [Answer](#)

Answer: (b) Root apex

Question 5.

Casparian strips occur in the cells of

- (a) Exodermis
- (b) Epiderms
- (c) Hypodermis
- (d) Endoderms

▼ [Answer](#)

Answer: (d) Endoderms

Question 6.

Lignified cells with narrow and pointed end wall are

- (a) Chlorenchyma
- (b) Parenchyma
- (c) Sclerenchyma
- (d) Endoderms

▼ [Answer](#)

Answer: (c) Sclerenchyma

Question 7.

Nucleus is absent in

- (a) Vessels
- (b) Sieve tube elements
- (c) Tracheid
- (d) All of these

▼ [Answer](#)

Answer: (d) All of these

Question 8.

Age of a tree is calculated by its

- (a) Girth
- (b) Height
- (c) Number of annual rings
- (d) Number of branches

▼ [Answer](#)

Answer: (c) Number of annual rings.

Question 9.

Youngest secondary xylem occurs

- (a) Just outside the vascular cambium
- (b) Just inside the vascular cambium
- (c) Just outside the vascular cambium
- (d) Just inside the cork cambium

▼ [Answer](#)

Answer: (b) Just inside the vascular cambium.

Question 10.

Mesophyll cells in a leaf are

- (a) Sclerenchymatous
- (b) Collenchymatous
- (c) Parenchymatous
- (d) Meristem

▼ [Answer](#)

Answer: (c) Parenchymatous

Question 11.

Healing of wounds occur due to the activity of

- (a) Intercalary meristem
- (b) Secondary meristem

- (c) Primary meristem
- (d) Apical meristem

▼ [Answer](#)

Answer: (b) Secondary meristem

Question 12.

Lateral root arise from

- (a) Cambium
- (b) Pericycle
- (c) Epidermis
- (d) Endodermis

▼ [Answer](#)

Answer: (b) Pericycle

Question 13.

Vascular bundles are absent in

- (a) Dicots
- (b) Monocots
- (c) Cambium
- (d) Pteridophytes

▼ [Answer](#)

Answer: (d) Pteridophytes

Question 14.

Which one contain only living cells?

- (a) Vessels
- (b) Sclerenchyma
- (c) Trachieds
- (d) Parenchyma

▼ [Answer](#)

Answer: (d) Parenchyma

Question 15.

Vascular bundle having cambium is

- (a) Closed
- (b) Open
- (c) Colleral
- (d) Conjoint

▼ [Answer](#)

Answer: (b) Open

Question 16.

Lignified cell wall occurs in

- (a) Xylem cells
- (b) Epidermal cells
- (c) Cambial cells
- (d) Phloem cells

▼ [Answer](#)

Answer: (a) Xylem cells .

Question 17.

Bordered pits are more common in

- (a) Gymnosperms
- (b) Monocots
- (c) Dicots
- (d) All of these

▼ [Answer](#)

Answer: (a) Gymnosperms

Question 18.

Both apical meristems and intercalary meristem are also called

- (a) Intercalary meristems
- (b) Lateral meristems
- (c) Primary meristems
- (d) Meristems

▼ [Answer](#)

Answer: (c) Primary meristems

Question 19.

Complex tissues also provides mechanical strength to the plant parts. It is composed elements

- (a) Trachieds
- (b) Vessels
- (c) Xylem fibres
- (d) Xylem parenchyma
- (e) All of these

▼ [Answer](#)

Answer: (e) All of these

Question 20.

The first formed xylem elements are called

- (a) Metaxylem
- (b) Endarch
- (c) Protoxylem
- (d) Exarch

▼ [Answer](#)

Answer: (c) Protoxylem

II. Fill in the blanks:

Question 1.

The plant is made up of Cells which are organised into and the tissues into

▼ [Answer](#)

Answer: tissues, organs

Question 2.

Plants have different kinds of

▼ [Answer](#)

Answer: meristems

Question 3.

Fascicular vascular cambium, interfascicular cambium and cork- cambium are examples of meristems.

▼ [Answer](#)

Answer: lateral

Question 4.

The various simple tissues are, and

▼ [Answer](#)

Answer: parenchyma, collenchyma, sclerenchyma

Question 5.

Parenchymatous cells form the major components like of and parts.

▼ [Answer](#)

Answer: cortex., pit, mesophyll, leaves, floral

Question 6.

..... forms below the epidermis, in dicotyle-donous plants.

▼ [Answer](#)

Answer: Collenchyma, hypodermis

Question 7.

..... consists of long, narrow cells with thick and lignified i cell walls having a few or numerous pits.

▼ [Answer](#)

Answer: Sclerenchyma

Question 8.

..... is cbmposed of four different kinds of elements, namely, trachieds, vessels, xylem fibres and xylem paranchyma.

▼ [Answer](#)

Answer: Complex tissues

Question 9.

..... are long, tube-like structures, arranged longitudinally and are associated with the companion cells.

▼ [Answer](#)

Answer: Sieve tube elements

Question 10.

Each stoma is composed of two bean-shaped cells known as the

▼ [Answer](#)

Answer: guard cells

Question 11.

The cells of epidermis bear a number of hair known as

▼ [Answer](#)

Answer: trichomes

Question 12.

The outermost layer is

▼ [Answer](#)

Answer: epidermis

Question 13.

The consists of several layers of thin-walled parenchymata with intercellular spaces.

▼ [Answer](#)

Answer: cortex

Question 14.

A number of are arranged in a ring. Each vascular bundle is, and

▼ [Answer](#)

Answer: vascular, conjoint, collateral endarch, open

Question 15.

When the cells in the leaves are turgid, the leaf surface is exposed.

▼ [Answer](#)

Answer: bulliform

III. Mark the statements True (T) or False (F)

Question 1.

The spring wood is lighter in colour and has a low density whereas the autumn wood is darker and has a higher density.

▼ [Answer](#)

Answer: True

Question 2.

The wood formed during spring season is called autumn wood or late wood.

▼ [Answer](#)

Answer: False

Question 3.

Phellogen, phellem, and phelloderm are collectively known as lenticel.

▼ [Answer](#)

Answer: False

Question 4.

The peripheral region of the secondary xylem, is lighter in colour and is known as the sapwood.

▼ [Answer](#)

Answer: True

Question 5.

Secondary growth also occur in stems and roots of gymnosperms. However, secondary growth does not occur in monocotyledons.

▼ [Answer](#)

Answer: True

Question 6.

All the dead cells lying outside the active cork cambium constitute the bark.

▼ [Answer](#)

Answer: True

Question 7.

Lenticels permit the exchange of gases between the outer at-mosphere and the internal tissue of the stem.

▼ [Answer](#)

Answer: True

Question 8.

The two kinds of woods appear as alternate concentric rings, constituting an annual ring.

▼ [Answer](#)

Answer: True

Question 9.

Vascular system includes vascular bundles, which can be seen in the veins.

▼ [Answer](#)

Answer: True

Question 10.

When xylem and phloem within a vascular bundle are arranged in an alternate manner on different radii as in the roots are called radial as in the roots.

▼ [Answer](#)

Answer: True

IV. Match the items or column I with the items of column II

Column I	Column II
(i) Heart wood	(a) spring wood
(ii) Spring season	(b) innermost layer of the cortex
(iii) Secondary growth	(c) unicellular
(iv) Isobilateral leaf	(d) tapering cylindrical cells

(v) Endodermis	(e) comprises dead elements with highly lignified walls.
(vi) Root hairs	(f) thick-walled, elongated and pointed cells.
(vii) Phloem parenchyma	(g) two lateral meristems vascular cambium and cork cambium.
(viii) Primary xylem	(h) they are mostly dead and without protoplasts
(ix) Xylem parenchymatous	(i) bulliform cells.
(x) Sclereids	(j) walls are made up of cellulose
(xi) Collenchyma	(k) hypodermis below the epidermis.
(xii) Parenchymatous	(l) cortex, pith, mesophyll of leaves, floral parts.
(xiii) Fibres	(m) protoxylem and metaxylem
(xiv) Sclerenchyma	(n) spherical, oval or cylindrical, highly thickened dead cells with very narrow cavities.

▼ Answer

Answer:

Column I	Column II
(i) Heart wood	(e) comprises dead elements with highly lignified walls.
(ii) Spring season	(a) spring wood
(iii) Secondary growth	(g) two lateral meristems vascular cambium and cork cambium.
(iv) Isobilateral leaf	(i) bulliform cells.
(v) Endodermis	(b) innermost layer of the cortex
(vi) Root hairs	(c) unicellular
(vii) Phloem parenchyma	(d) tapering cylindrical cells
(viii) Primary xylem	(m) protoxylem and metaxylem
(ix) Xylem parenchymatous	(j) walls are made up of cellulose
(x) Sclereids	(n) spherical, oval or cylindrical, highly thickened dead cells with very narrow cavities.
(xi) Collenchyma	(k) hypodermis below the epidermis.
(xii) Parenchymatous	(l) cortex, pith, mesophyll of leaves, floral parts.
(xiii) Fibres	(f) thick-walled, elongated and pointed cells.
(xiv) Sclerenchyma	(h) they are mostly dead and without protoplasts