

Relations and Functions

Question 1.

The domain of the function ${}^{7-x}P_{x-3}$ is

- (a) $\{1, 2, 3\}$
- (b) $\{3, 4, 5, 6\}$
- (c) $\{3, 4, 5\}$
- (d) $\{1, 2, 3, 4, 5\}$

Answer: (c) $\{3, 4, 5\}$

The function $f(x) = {}^{7-x}P_{x-3}$ is defined only if x is an integer satisfying the following inequalities:

- 1. $7 - x \geq 0$
- 2. $x - 3 \geq 0$
- 3. $7 - x \geq x - 3$

Now, from 1, we get $x \leq 7$ 4

from 2, we get $x \geq 3$ 5

and from 3, we get $x \leq 5$ 6

From 4, 5 and 6, we get

$$3 \leq x \leq 5$$

So, the domain is $\{3, 4, 5\}$

Question 2.

The domain of $\tan^{-1} (2x + 1)$ is

- (a) $\mathbb{R}$
- (b) $\mathbb{R} - \{1/2\}$
- (c) $\mathbb{R} - \{-1/2\}$
- (d) None of these

Answer: (a) $\mathbb{R}$

Since $\tan^{-1} x$ exists if $x \in (-\infty, \infty)$

So, $\tan^{-1} (2x + 1)$ is defined if

$$-\infty < 2x + 1 < \infty$$

$$\Rightarrow -\infty < x < \infty$$

$$\Rightarrow x \in (-\infty, \infty)$$

$$\Rightarrow x \in \mathbb{R}$$

So, domain of $\tan^{-1}(2x + 1)$ is $\mathbb{R}$.

Question 3.

Two functions f and g are said to be equal if f

- (a) the domain of f = the domain of g
- (b) the co-domain of f = the co-domain of g
- (c) $f(x) = g(x)$ for all x
- (d) all of above

Answer: (d) all of above

Two functions f and g are said to be equal if f

- 1. the domain of f = the domain of g
 - 2. the co-domain of f = the co-domain of g
 - 3. $f(x) = g(x)$ for all x
-

Question 4.

If the function $f: \mathbb{R} \rightarrow \mathbb{R}$ be given by $f(x) = x^2 + 2$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ is given by $g(x) = x/(x - 1)$. The value of $\text{gof}(x)$ is

- (a) $(x^2 + 2)/(x^2 + 1)$
- (b) $x^2/(x^2 + 1)$
- (c) $x^2/(x^2 + 2)$
- (d) none of these

Answer: (a) $(x^2 + 2)/(x^2 + 1)$

Given $f(x) = x^2 + 2$ and $g(x) = x/(x - 1)$

Now, $\text{gof}(x) = g(x^2 + 2) = (x^2 + 2)/(x^2 + 2 - 1) = (x^2 + 2)/(x^2 + 1)$

Question 5.

Given $g(1) = 1$ and $g(2) = 3$. If $g(x)$ is described by the formula $g(x) = ax + b$, then the value of a and b is

- (a) 2, 1
- (b) -2, 1
- (c) 2, -1
- (d) -2, -1

Answer: (c) 2, -1

Given, $g(x) = ax + b$

Again, $g(1) = 1$

$$\Rightarrow a \times 1 + b = 1$$

$$\Rightarrow a + b = 1 \dots\dots\dots 1$$

$$\text{and } g(2) = 3$$

$$\Rightarrow a \times 2 + b = 3$$

$$\Rightarrow 2a + b = 3 \dots\dots\dots 2$$

Solve equation 1 and 2, we get

$$a = 2, b = -1$$

Question 6.

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function given by $f(x) = x^2 + 1$ then the value of $f^{-1}(26)$ is

(a) 5

(b) -5

(c) ± 5

(d) None of these

Answer: (c) ± 5

$$\text{Let } y = f(x) = x^2 + 1$$

$$\Rightarrow y = x^2 + 1$$

$$\Rightarrow y - 1 = x^2$$

$$\Rightarrow x = \pm\sqrt{y - 1}$$

$$\Rightarrow f^{-1}(x) = \pm\sqrt{x - 1}$$

$$\text{Now, } f^{-1}(26) = \pm\sqrt{26 - 1}$$

$$\Rightarrow f^{-1}(26) = \pm\sqrt{25}$$

$$\Rightarrow f^{-1}(26) = \pm 5$$

Question 7.

the function $f(x) = x - [x]$ has period of

(a) 0

(b) 1

(c) 2

(d) 3

Answer: (b) 1

Let T is a positive real number.

Let $f(x)$ is periodic with period T .

Now, $f(x + T) = f(x)$, for all $x \in \mathbb{R}$

$$\Rightarrow x + T - [x + T] = x - [x], \text{ for all } x \in \mathbb{R}$$

$$\Rightarrow [x + T] - [x] = T, \text{ for all } x \in \mathbb{R}$$

Thus, there exist $T > 0$ such that $f(x + T) = f(x)$ for all $x \in \mathbb{R}$

Now, the smallest value of T satisfying $f(x + T) = f(x)$ for all $x \in \mathbb{R}$ is 1
So, $f(x) = x - [x]$ has period 1

Question 8.

The function $f(x) = \sin(\pi x/2) + \cos(\pi x/2)$ is periodic with period

- (a) 4
- (b) 6
- (c) 12
- (d) 24

Answer: (a) 4

Period of $\sin(\pi x/2) = 2\pi/(\pi/2) = 4$

Period of $\cos(\pi x/2) = 2\pi/(\pi/2) = 4$

So, period of $f(x) = \text{LCM}(4, 4) = 4$

Question 9.

The domain of the function $f(x) = x/(1 + x^2)$ is

- (a) $\mathbb{R} - \{1\}$
- (b) $\mathbb{R} - \{-1\}$
- (c) $\mathbb{R}$
- (d) None of these

Answer: (c) $\mathbb{R}$

Given, function $f(x) = x/(1 + x^2)$

Since $f(x)$ is defined for all real values of x .

So, $\text{domain}(f) = \mathbb{R}$

Question 10.

If $f : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = x^2 - 3x + 2$, the $f(f(y))$ is

- (a) $x^4 + 6x^3 + 10x^2 + 3x$
- (b) $x^4 - 6x^3 + 10x^2 + 3x$
- (c) $x^4 + 6x^3 + 10x^2 - 3x$
- (d) $x^4 - 6x^3 + 10x^2 - 3x$

Answer: (d) $x^4 - 6x^3 + 10x^2 - 3x$

Given, $f(x) = x^2 - 3x + 2$

Now, $f(f(y)) = f(x^2 - 3x + 2)$

$= (x^2 - 3x + 2)^2 - 3(x^2 - 3x + 2) + 2$

$= x^4 - 6x^3 + 10x^2 - 3x$

Question 11.

If n is the smallest natural number such that $n + 2n + 3n + \dots + 99n$ is a perfect square, then the number of digits in square of n is

- (a) 1
- (b) 2
- (c) 3
- (d) 4

Answer: (c) 3

Given that

$$n + 2n + 3n + \dots + 99n$$

$$= n \times (1 + 2 + 3 + \dots + 99)$$

$$= (n \times 99 \times 100)/2$$

$$= n \times 99 \times 50$$

$$= n \times 9 \times 11 \times 2 \times 25$$

To make it perfect square we need 2×11

$$\text{So } n = 2 \times 11 = 22$$

$$\text{Now } n^2 = 22 \times 22 = 484$$

So, the number of digit in $n^2 = 3$

Question 12.

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function defined by $f(x) = \cos(5x + 2)$, then f is

- (a) injective
- (b) surjective
- (c) bijective
- (d) None of these

Answer: (d) None of these

$$\text{Given, } f(x) = \cos(2x + 5)$$

$$\text{Period of } f(x) = 2\pi/5$$

Since $f(x)$ is a periodic function with period $2\pi/5$, so it is not injective.

The function f is not surjective also as its range $[-1, 1]$ is a proper subset of its co-domain $\mathbb{R}$

Question 13.

The function $f(x) = \sin(\pi x/2) + 2\cos(\pi x/3) - \tan(\pi x/4)$ is periodic with period

- (a) 4
- (b) 6
- (c) 8
- (d) 12

Answer: (d) 12

Period of $\sin(\pi x/2) = 2\pi/(\pi/2) = 4$

Period of $\cos(\pi x/3) = 2\pi/(\pi/3) = 6$

Period of $\tan(\pi x/4) = \pi/(\pi/4) = 4$

So, period of $f(x) = \text{LCM}(4, 6, 4) = 12$

Question 14.

If the function $f: \mathbb{R} \rightarrow \mathbb{R}$ be given by $f(x) = x^2 + 2$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ is given by $g(x) = x/(x-1)$. The value of $\text{gof}(x)$ is

(a) $(x^2 + 2)/(x^2 + 1)$

(b) $x^2/(x^2 + 1)$

(c) $x^2/(x^2 + 2)$

(d) none of these

Answer: (a) $(x^2 + 2)/(x^2 + 1)$

Given $f(x) = x^2 + 2$ and $g(x) = x/(x-1)$

Now, $\text{gof}(x) = g(x^2 + 2) = (x^2 + 2)/(x^2 + 2 - 1) = (x^2 + 2)/(x^2 + 1)$

Question 15.

The domain of the function ${}^{7-x}P_{x-3}$ is

(a) $\{1, 2, 3\}$

(b) $\{3, 4, 5, 6\}$

(c) $\{3, 4, 5\}$

(d) $\{1, 2, 3, 4, 5\}$

Answer: (c) $\{3, 4, 5\}$

The function $f(x) = {}^{7-x}P_{x-3}$ is defined only if x is an integer satisfying the following inequalities:

1. $7 - x \geq 0$

2. $x - 3 \geq 0$

3. $7 - x \geq x - 3$

Now, from 1, we get $x \leq 7$ 4

from 2, we get $x \geq 3$ 5

and from 3, we get $x \leq 5$ 6

From 4, 5 and 6, we get

$3 \leq x \leq 5$

So, the domain is $\{3, 4, 5\}$

Question 16.

If $f(x) = e^x$ and $g(x) = \log_e x$ then the value of $\text{fog}(1)$ is

- (a) 0
- (b) 1
- (c) -1
- (d) None of these

Answer: (b) 1

Given, $f(x) = e^x$

and $g(x) = \log x$

$\text{fog}(x) = f(g(x))$

$= f(\log x)$

$= e^{\log x}$

$= x$

So, $\text{fog}(1) = 1$

Question 17.

A relation R is defined from the set of integers to the set of real numbers as $(x, y) \in R$ if $x^2 + y^2 = 16$ then the domain of R is

- (a) (0, 4, 4)
- (b) (0, -4, 4)
- (c) (0, -4, -4)
- (d) None of these

Answer: (b) (0, -4, 4)

Given that:

$(x, y) \in R \Leftrightarrow x^2 + y^2 = 16$

$\Leftrightarrow y = \pm\sqrt{16 - x^2}$

when $x = 0 \Rightarrow y = \pm 4$

$(0, 4) \in R$ and $(0, -4) \in R$

when $x = \pm 4 \Rightarrow y = 0$

$(4, 0) \in R$ and $(-4, 0) \in R$

Now for other integral values of x, y is not an integer.

Hence $R = \{(0, 4), (0, -4), (4, 0), (-4, 0)\}$

So, $\text{Domain}(R) = \{0, -4, 4\}$

Question 18.

The period of the function $f(x) = \sin(2\pi x/3) + \cos(\pi x/3)$

- (a) 3
- (b) 4
- (c) 12
- (d) None of these

Answer: (c) 12

Given, function $f(x) = \sin(2\pi x/3) + \cos(\pi x/2)$

Now, period of $\sin(2\pi x/3) = 2\pi/\{(2\pi/3)\} = (2\pi \times 3)/(2\pi) = 3$

and period of $\cos(\pi x/2) = 2\pi/\{(\pi/2)\} = (2\pi \times 2)/(\pi) = 2 \times 2 = 4$

Now, period of $f(x) = \text{LCM}(3, 4) = 12$

Hence, period of function $f(x) = \sin(2\pi x/3) + \cos(\pi x/2)$ is 12

Question 19.

If $f(x) = ax + b$ and $g(x) = cx + d$ and $f\{g(x)\} = g\{f(x)\}$ then

(a) $f(a) = g(c)$

(b) $f(b) = g(b)$

(c) $f(d) = g(b)$

(d) $f(c) = g(a)$

Answer: (c) $f(d) = g(b)$

Given, $f(x) = ax + b$ and $g(x) = cx + d$ and

Now, $f\{g(x)\} = g\{f(x)\}$

$\Rightarrow f\{cx + d\} = g\{ax + b\}$

$\Rightarrow a(cx + d) + b = c(ax + b) + d$

$\Rightarrow ad + b = cb + d$

$\Rightarrow f(d) = g(b)$

Question 20.

The domain of the function $f(x) = 1/(2 - \cos 3x)$ is

(a) $(1/3, 1)$

(b) $[1/3, 1)$

(c) $(1/3, 1]$

(d) $\mathbb{R}$

Answer: (d) $\mathbb{R}$

Given

function is $f(x) = 1/(2 - \cos 3x)$

Since $-1 \leq \cos 3x \leq 1$ for all $x \in \mathbb{R}$

So, $-1 \leq 2 - \cos 3x \leq 1$ for all $x \in \mathbb{R}$

$\Rightarrow f(x)$ is defined for all $x \in \mathbb{R}$

So, domain of $f(x)$ is $\mathbb{R}$
