




BANGALORE SAHODAYA SCHOOLS COMPLEX ASSOCIATION
QUESTION PAPER (2023-24) PHYSICS (Code – 042)
MARKING SCHEME CLASS XII – SET 1


Q.NO	SECTION A	Marks
1	(c) $F = F' \text{ or } \frac{Q_1 Q_2}{4\pi\epsilon_0 r^2} = \frac{Q_1 Q_2}{4\pi\epsilon_0 (r')^2 k}$ $r' = r/\sqrt{k}$	1
2	(a) The electric field $E = -dV/dx$ $\Rightarrow E = 16x - 4$. The electric field is along x-direction. As the electric field is always perpendicular to the equipotential surface, the equipotential surface must be planes parallel to y-z plane.	1
3	(c) $i_A = 2A, r_A = 2cm, \theta_A = 2\pi - \frac{\pi}{2} = \frac{3\pi}{2}$ $i_B = 3A, r_B = 4cm, \theta_B = 2\pi - \frac{\pi}{3} = \frac{5\pi}{3}$ $B = \frac{\mu_0 I \theta}{4\pi R}$ $B_A = i_A \times \frac{\theta_A R_B}{i_B \theta_B R_A} = 6$ $B_B = \frac{i_B \theta_B R_A}{i_A \theta_A R_B} = 5$	1
4	(b) The net magnetic force on a current carrying closed loop is zero. Here, $\vec{F}_{BC} = \vec{F}$ $\vec{F}_{AB} = 0$ $\therefore \vec{F}_{AB} + \vec{F}_{BC} + \vec{F}_{AC} = 0$ $\Rightarrow \vec{F}_{AC} = -\vec{F}_{BC} \quad (\vec{F}_{AB} = 0)$	1
5	(c) Dipole moment of circular loop is m . $m_1 = IA = I\pi R^2 \{R = \text{radius of the loop}\}$ $B_1 = \frac{\mu_0 I}{2R}$ moment becomes double $\Rightarrow R$ becomes $\sqrt{2}R$ (keeping current constant) $m_2 = I\pi (\sqrt{2}R)^2 = 2I\pi R^2 = 2m_1$ $B_2 = \frac{\mu_0 I}{2(\sqrt{2}R)} = \frac{B_1}{\sqrt{2}}$ $B_1 = \frac{B_2}{\sqrt{2}}$ $B_2 = \frac{B_1}{\sqrt{2}}$	1
6	(a) For a solenoid of n turns per unit length carrying current I , $H = nI$. $\therefore M = (\mu_r - 1)nI$ $M = (1000 - 1) \times 1000 \times 0.5$	1


	$M = 5 \times 10^5 \text{ Am}^{-1}$ As magnetic moment, $m = M \times V$ $\therefore m = 5 \times 10^5 \times 10^{-3} = 500 \text{ Am}^2$	
7	(c) Given $n = 2 \times 10^4$; $I = 4 \text{ A}$ Initially, the magnetic field at the centre of the solenoid is given as $B_i = \mu_0 n I = 4\pi \times 10^{-7} \times 2 \times 10^4 \times 4 = 32\pi \times 10^{-3} \text{ T}$ Initial magnetic flux through the coil is given as $\phi_i = NBA = 100 \times 32\pi \times 10^{-3} \times \pi \times (0.01)^2 = \phi_i = 32\pi^2 \times 10^{-5} \text{ Tm}^2$ Finally $I = 0 \text{ A}$ $\therefore B_f = 0$ or $\phi_f = 0$ Induced charge, $q = \Delta\phi /R = \phi_f - \phi_i /R = 32\pi^2 \times 10^{-5}/10\pi^2 = 32 \times 10^{-6} \text{ C} = 32 \mu\text{C}$	1
8	(d) The phase difference ϕ between current and voltage is given by $\tan\phi = \frac{X_C - X_L}{R}$ Or $\frac{X_C - X_L}{R} = \tan 45^\circ = 1$ Or $X_C = X_L + R$ Or $\frac{1}{2\pi f C} = 2\pi f L + R$ Or $C = \frac{1}{2\pi f (2\pi f L + R)}$	1
9	(a)	1
10	(b) $K = \frac{hc}{\lambda} - \phi$ and that in the second case is $K_{\text{max}_2} = \frac{hc}{\lambda} - \phi_0 = \frac{2hc}{\lambda} - \phi_0$ But $K_{\text{max}_2} = 3K_{\text{max}_1}$ (given) $\therefore \frac{2hc}{\lambda} - \phi_0 = 3 \left(\frac{hc}{\lambda} - \phi_0 \right)$ $\frac{2hc}{\lambda} - \phi_0 = \frac{3hc}{\lambda} - 3\phi_0$	1


	$3\phi_0 - \phi_0 = \frac{3hc}{\lambda} - \frac{2hc}{\lambda}$ $2\phi_0 = \frac{hc}{\lambda} \text{ or } \phi_0 = \frac{hc}{2\lambda}$	
11	<p>(a)</p> <p>Explanation: Kinetic energy of any charge q accelerated by V volt, $K = qV$</p> <p>$\Rightarrow K_\alpha = 2eV, \quad K_p = eV$</p> <p>At distance of closest approach, $K = U$</p> <p>Let atomic number of target nucleus be Z</p> <p>For α-particle, $2eV = \frac{2Ze^2}{4\pi\epsilon_0 r}$</p> <p>For proton, $eV = \frac{Ze^2}{4\pi\epsilon_0 r'}$</p> <p>Hence $r' = r$</p>	1
12	<p>(c)</p> <p>For Lyman Series wavelength will be longest when the electron has transition from $n=2$ to $n=1$ level.</p> <p>$\Rightarrow \frac{1}{\lambda} = R \left(\frac{1}{1^2} - \frac{1}{2^2} \right)$</p> <p>For Balmer Series wavelength will be longest when the electron has transition from $n=3$ to $n=2$ level.</p> <p>$\Rightarrow \frac{1}{\lambda_B} = R \left(\frac{1}{2^2} - \frac{1}{3^2} \right)$</p> <p>Hence, $\frac{\lambda}{\lambda_B} = \frac{5}{27}$</p>	1
13	(a)	1
14	(b)	1
15	(c)	1
16	(c)	1
17	<p>Resistance of heaters $R_1 = 400/3 \, \Omega$, $R_2 = 400/6 \, \Omega$</p> <p>When heaters are connected in series, current in circuit,</p> $I = \frac{V}{R_1 + R_2} = \frac{200}{\frac{400}{3} + \frac{400}{6}} = 1A$ <p>Heat produced in 200V, 300 W heater per second</p> $Q_1 = I^2 R_1 = (1)^2 \times \frac{400}{3} = 133.33 \text{ J s}^{-1}$ <p>Heat produced in 200 V and 600 W heater per second.</p>	1

	$Q_2 = I^2 R_2 = (1)^2 \times \frac{400}{6} = 66.66 \text{ Js}^{-1}$ <p>Clearly heat produced in 300 W heater is more than that produced in 600 W heater</p>	1
18	<p>For total internal reflection at the vertical face, $i = i_c$ (if μ is minimum)</p> <p>If $r =$ angle of refraction of ray into the prism</p> <p>Clearly, $r + i_c = 90^\circ$</p> <p>$r = 90^\circ - i_c$</p> $\mu = \frac{\sin i}{\sin r} = \frac{\sin 45^\circ}{\sin(90-i_c)} = \frac{1}{\sqrt{2} \cos i_c}$ <p>Also, $\mu = \frac{\sin i_c}{\sin i} \Rightarrow \sin i = \sqrt{2} \cos i_c$</p> <p>Or, $\tan i_c = \sqrt{2}$</p> <p>Hence $\sin i_c = \frac{\sqrt{2}}{\sqrt{3}}$</p> $\mu = \frac{1}{\sin i_c} = \frac{\sqrt{3}}{2}$	<p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
19	<p>Let phase difference be ϕ</p> $I = 4I_0 \cos^2(\phi/2) = 4I_0/2 = 2I_0$ <p>$\Rightarrow \phi = \pi/2$</p> <p>Path difference $= \phi \lambda / 2\pi = \lambda/4$</p> <p>$\Rightarrow y = \lambda D / 4d$</p> <p>OR</p> <p>For λ_1, $\delta_m = A$, $n_1 = \sqrt{3}$</p> <p>For λ_2, $\delta_m = 30^\circ$, $n_2 = ?$</p> <p>Hence, $n_1 = \sqrt{3} = \frac{\sin(\frac{A+A}{2})}{\sin(\frac{A}{2})} = \frac{\sin A}{\sin \frac{A}{2}} = \frac{2 \sin \frac{A}{2} \cos \frac{A}{2}}{\sin \frac{A}{2}} = 2 \cos \frac{A}{2}$</p> $\frac{\sqrt{3}}{2} = \cos \frac{A}{2} \Rightarrow A = 60^\circ$ $n_2 = \frac{\sin(\frac{60+30}{2})}{\sin 30} = \frac{\sin 45}{\sin 30} = \sqrt{2}$	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>1</p>
20	<p>(i) $\frac{\text{Energy of photon}}{\text{K.E. of electron}} = \frac{2m\lambda c}{h}$</p> $= \frac{2 \times 9.11 \times 10^{-31} \times 10^{-9} \times 3 \times 10^8}{6.6 \times 10^{-34}}$ $= \frac{9110}{11} = 9110:11$ <p>(ii) Any two features that cannot be explained by wave theory of light</p>	<p>1</p> <p>1</p>

21	 <p>Explanation</p>	1 1
22	<p>Equivalent capacitance = $(200/3)\mu\text{F}$</p> <p>Voltage across $C_1 = 100\text{V}$</p> <p>Voltage across $C_2 = 50\text{V}$</p> <p>Charge across $C_1 = \text{Charge across } C_2 = 10^{-8}\text{ C}$</p>	1 $\frac{1}{2}$ $\frac{1}{2}$ 1
23	<p>The three cells are in parallel and hence effective emf = $40/3\text{ V}$</p> <p>There will be no current in the branch having capacitor after complete charging</p> <p>Hence the charge on the capacitor $q = CV = 200/3\text{ }\mu\text{C}$</p>	2 1
24	<p>(a) Diagram</p> <p>Expression and Direction of B</p> <p>Expression and Direction of force per unit length</p> <p>Definition of ampere</p> <p>Or</p> <p>(b) (i) By connecting a small resistance in parallel with the galvanometer</p> <p>Derivation of S</p> <p>Effective resistance</p> <p>(ii) $I_s = \frac{\theta}{I}, V_s = \frac{I_s}{R}$</p> $I'_s = I_s + \frac{50}{100} I_s = \frac{3}{2} I_s$ $V'_s = \frac{I'_s}{2R} = \frac{\frac{3}{2} I_s}{2R} = \frac{3}{4} V_s$ <p>$V'_s = 0.75\% V_s$</p> <p>So, V_s decreases by 25%.</p>	$\frac{1}{2}$ 1 1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

25	<p>Diagram</p> <p>Derivation</p> <p>$M_{12} = M_{21}$</p>	<p>1</p> <p>1</p> <p>1</p>
26	<p>(a) $\lambda_1 \rightarrow \text{IR}, \lambda_2 \rightarrow \text{radiowaves } \lambda_3 \rightarrow \text{X-rays}$</p> <p>$\lambda_2 > \lambda_1 > \lambda_3$</p> <p>(b) Figure</p>	<p>$1\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p>
27	<p>(a) Statement of Bohr's postulate of quantisation of angular momentum</p> <p>Justification using de Broglie hypothesis</p> <p>(b) For third excited state, $n=4$</p> <p>Now, the total number of possible spectral lines is given by the formula,</p> <p>$N=n(n-1)/2$</p> <p>On putting $n=4$</p> <p>$\Rightarrow N=4(4-1)/2$</p> <p>Hence, we get</p> <p>$N=6$</p>	<p>$\frac{1}{2}$</p> <p>$1\frac{1}{2}$</p> <p>1</p>
28	 <p>Binding energy per nucleon of lighter nuclei is small. In an attempt to get higher B.E/A, lighter nuclei undergo nuclear fusion.</p> <p>Saturation effect / short range nature of nuclear force.</p>	<p>$1\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p>
29	<p>(i)(b) Eyepiece acts as a simple microscope. It forms a virtual and erect final image.</p> <p>$f_e = 6.25 \text{ cm}, v_e = -25 \text{ cm}$</p> <p>using $\frac{1}{f_e} = \frac{1}{v_e} - \frac{1}{u_e}$</p> <p>we get, $u_e = -5 \text{ cm}$</p> <p>(ii)(b)</p>	<p>1</p>

	$L = v_0 + u_e \Rightarrow v_0 = L - u_e $ $\Rightarrow v_0 = 15 - 5 = 10 \text{ cm}$ $f_0 = 2 \text{ cm}$ $\frac{1}{f_o} = \frac{1}{v_o} - \frac{1}{u_o}$ <p>Substituting, we get $u_o = -2.5 \text{ cm}$</p> $(iii)(a) M = \frac{v_0}{u_0} \left(1 + \frac{D}{f_e}\right) = \frac{10}{2.5} \left(1 + \frac{25}{6.25}\right) = 20$ <p>(iv) (a)</p> <p>OR</p> <p>(d)</p>	<p>1</p> <p>1</p> <p>1</p>
30	<p>i) (c) Doping increases the resistivity of semiconductor</p> <p>OR</p> <p>Which of the following energy band diagram shows the n type semiconductor?</p> <p>(d)</p>  <p>ii) (b) $E_{\min} = \frac{hc}{\lambda_{\max}}$</p> $\lambda_{\max} = 589 \text{ nm}$ <p>iii) (a) Width of the depletion region,</p> $d = V/E = 0.4/10^6 = 4 \times 10^{-7} \text{ m}$ <p>iv) (c) The equivalent circuit is</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p>




After observing this circuit we get that,

$$i = 10/2 = 5 \text{ mA} = i_2$$

$$i_1 = 0 \text{ mA}$$

31

a) (i)



The electric field at the point P due to +q placed at B is,

$$E_1 = \frac{1}{4\pi\epsilon_0} \frac{q}{(r-d)^2} \text{ (along BP)}$$

The electric field at the point P due to -q placed at A is,

$$E_2 = \frac{1}{4\pi\epsilon_0} \frac{q}{(r+d)^2} \text{ (along PA)}$$

Therefore, the magnitude of resultant electric field (E) acts in the direction of the vector with a greater, magnitude. The resultant electric field at P is


$$E = E_1 + (-E_2)$$

$$E = \frac{q}{4\pi\epsilon_0} \left[\frac{4rd}{(r^2 - d^2)^2} \right] \text{ along BP}$$

If the point P is far away from the dipole, then $d \ll r$

$$E = \frac{1}{4\pi\epsilon_0} \frac{2p}{r^3} \text{ along BP}$$

ii) Two point charges q and -q are located at point (0,0,-a) and (0,0,a).



(1) The electrostatic potential at (0, 0, z)

$$V = \frac{1}{4\pi\epsilon_0} \frac{-q}{AP} + \frac{1}{4\pi\epsilon_0} \frac{+q}{BP} = \frac{P}{4\pi\epsilon_0(z^2 - a^2)}$$

and The electrostatic potential at (x, y, 0)

$$V = \frac{1}{4\pi\epsilon_0} \frac{-q}{AQ} + \frac{1}{4\pi\epsilon_0} \frac{+q}{BQ}$$

Since, AQ=BQ
We have, electric potential at (x,y,0)=0.

(2)

Potential at (5, 0, 0)

$$V_1 = \frac{-q}{4\pi\epsilon_0} \frac{1}{\sqrt{(5-0)^2 + (-a)^2}} + \frac{q}{4\pi\epsilon_0} \frac{1}{\sqrt{(5-0)^2 + a^2}} = \frac{-q}{4\pi\epsilon_0} \sqrt{25 + a^2} = 0$$

Potential at point (-7, 0, 0)


$$V_2 = \frac{-q}{4\pi\epsilon_0} \frac{1}{\sqrt{(-7-0)^2 + a^2}} + \frac{q}{4\pi\epsilon_0} \frac{1}{\sqrt{(-7-0)^2 + a^2}} = \frac{-q}{4\pi\epsilon_0} \cdot \frac{1}{\sqrt{4a + a^2}} + \frac{q}{4\pi\epsilon_0} \frac{1}{\sqrt{4a + a^2}} = 0$$

$$\text{Work done} = \text{Charge} \times \text{Potential} \left(V = \frac{W}{Q} \right)$$

$$\text{Charge} \times (V_2 - V_1) = \text{Charge} \times 0 - 0 \\ = 0 \quad W = 0$$

OR

(b) (i)



Let σ be the surface charge density of the sheet. From symmetry, E on either side of the sheet must be perpendicular to the plane of the sheet having same magnitude at all points equidistant from the sheet.

We take a cylinder of cross-sectional area A and length $2r$ as the Gaussian surface.

$$\text{Net flux through the flat surface} = EA + EA = 2EA$$

The flux through curved surface are zero because E and dA are at right angle,


\therefore Total electric flux over the entire surface of cylinder

$$\phi_E = 2EA$$

$$\text{Total charge enclosed by the cylinder } q = \sigma A$$

$$\text{According to Gauss's law } \phi_E = q/\epsilon_0$$

$$\therefore 2EA = \sigma A/\epsilon_0 \text{ or } E = \sigma/2\epsilon_0$$

	<p>ii) When no electric field is applied, the time period of oscillation is:</p> $T = 2\pi\sqrt{\frac{l}{g}}$ <p>When electric field is applied, T'</p> $= 2\pi\sqrt{\frac{l}{g-a}} \quad [a = \frac{qE}{m} = 2.5]$ <p>solving above two equations T'=2.6 s</p> <p>Therefore time taken for 25 oscillations = 25 T' = 65 s</p>	<p>1</p> <p>1</p>
32	<p>a) (i) X → capacitor</p> <p>(ii) curve B → voltage</p> <p>curve C → current</p> <p>curve A → power</p> <p>(iii) $X_c = \frac{1}{\omega C}$</p> <p>$X_c \propto \frac{1}{\omega}$</p>  <p>(iv) Derivation of expression for the current</p> <p>Phase relation</p> <p style="text-align: center;">OR</p> <p>(b) (i) Diagram</p> <p>Principle</p> <p>Working</p> <p>(ii) $I_s = \frac{E_s}{R_s} = \frac{22}{440} = \frac{1}{20} \text{ A}$</p> $\eta = \frac{E_s I_s}{E_p I_p}$ $\frac{90}{100} = \frac{22 \times \left(\frac{1}{20}\right)}{220 \times I_p}$ $\Rightarrow I_p = 0.0056 \text{ A}$ <p>(iii) There is no change in magnetic field due to dc</p>	<p>1/2</p> <p>1 1/2</p> <p>1/2</p> <p>1/2</p> <p>1 1/2</p> <p>1/2</p> <p style="text-align: center;">OR</p> <p>1</p> <p>1/2</p> <p>1 1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>
33	<p>(a) (i) Ray diagram</p> <p>Derivation of $\frac{n_2}{v} - \frac{n_1}{u} = \frac{n_2 - n_1}{R}$</p>	<p>1</p> <p>2</p> <p>1</p>

